

Review Article

PAPAYA (*CARICA PAPAYA*): SOURCE MATERIAL FOR ANTICANCER

***Sheikh Fauziya and R Krishnamurthy**

*CG Bhakta Institute of Biotechnology, Uka Tarsadia University, Maliba Campus Bardoli,
Dist. Surat, Gujarat, India-394350*

**Author for Correspondence*

ABSTRACT

Papaya (*Carica papaya* Linn.) is well known for its exceptional nutritional and medicinal properties throughout the world. The whole plant including its leaves, fruits, seed, bark, latex and their juice are use as medicine. Papaya is considering as Nutraceutical fruits duo its multifaceted properties. The whole plant of papaya contain enzyme Papain, Lycopene, Isothyocynate, important Mineral, Vitamins, Carbohydrates, Carotenoids, Flavonoids. These important nutritious fruits feed the body and immune system. A papaya in vitro study shows that it will treat many cancer cell line and papaya physiochemical having anticancer activities. Papaya is rich in enzyme papain which is effective against cancer. Papain breaks down the fibrin cancer cell wall and protein into amino acid form. Other than papain it also contain lycopene which highly reactive towards oxygen and free radical. Isothyocynate effective against breast, lung, colon pancreas, prostate as well as leukemia. These enzymes capable of inhibiting both formation and development of cancer cell.

Key Words: *Carica papaya, Anticancer, Physiochemical, Papain, Lycopene, Isothyocynate*

INTRODUCTION

The papaya tree is belonging to small family caricaceae having four genera in world. The genus carica linn is represented by four species in India, of which *Carica papaya* linn. is most widely cultivated and best-known species (Jean *et al.*, 2001). Papaya belongs to *Caricaceae* family. Scientific name of papaya is *Carica papaya*. It is commonly known as Papaya Melon tree, Pawpaw or Papau, Kapaya, Lapaya, Papyas, Papye, Tapayas, Fan mu gua (Bhattachrjee, 2001).

The fruit, leaves and latex obtain from papaya plant used medicinally and for various other purposes. Papain, major chemical compound extracted from fruit and stem latex is used in brewing and wine making and in textile and tanning industries (CSIR, 1992).

According to the Journal of Ethno pharmacology published on the 17th of February, 2010, international doctors and researchers from US and Japan have discovered that enzymes found in Papaya Leaf Tea have dramatic cancer-fighting properties against a broad range of tumors (Mike, 2012).

A study conducted by University of Florida researchers Nam Dang and colleagues in Japan has documented papaya's powerful anticancer properties and impact numerous lab-grown-tumors.

Papaya is small, erect, usually unbranched, fast-growing tree growing 3-6 meters high. Trunk is soft and grayish, marked with large petiole-scars. Leaves are somewhat rounded in outline, 1 meter broad or less, palmately 7-or 9-loabrd, each lobe pinnatelyincised or lobed. Petioles are stout, hollow, and about 1 meter long. Male flowers are in crowded cluster, straw-colored, and fragrant. Corolla tube is slander, about 2cm long. Female flowers are in short, axillary spikes or racemes, the petal 7cm long or less. Fruit is Indehiscent, subglobose, obovoid or oblong-cylindric, 5-30cm long, fleshy and yellowish or yellow-orange when ripe, containing numerous black seeds which are embedded in sweet pulp (Nadkarni, 1954).

MATERIALS AND METHODS

Origin and Distribution

Papaya is probably originated in southern Mexico and Costa Rica, subsequently it was crop introduce in Australia, Hawaii, Philippines, Srilanka, Southafrica, India and all tropical and subtropical regions. It is growing both commercially and in home garden. (Marotta *et al.*, 2006).

Review Article

Table 1: Nutritional value

Constituents	Ripe Papaya	Green Papaya
Water	89.1	92.6
Protein	8.26 g	10.8 g
Total lipid	0.93 g	1.35 g
Ash	4.59 g	6.76 g
Carbohydrates	86.2 g	81.1 g
Total DF	11.9 g	27.0 g
Sugar	NR	NR
Mineral		
Macronutrient		
Sodium	128.4 mg	283.8 mg
Potassium	1238 mg	2743 mg
Magnesium	229.4 mg	635.1 mg
Phosphorus	NR	NR
Calcium	146.8 mg	432.4 mg
Micronutrient		
Iron	12.84 mg	8.11 mg
Copper	0.18 mg	0.14 mg
Zinc	0.92 mg	0
Manganese	NR	NR
Selenium	NR	NR
Vitamins		
Vitamin C,	568.8 mg	391.9 mg
Thiamin	0.28 mg	0.54 mg
Riboflavin	0.28 mg	26 mg
Niacin	2.80 mg	4.05 mg
Pantothenic Acid	NR	NR
Vitamin B-6	NR	NR
Total Folate,	NR	NR
Folate	NR	NR
Vitamin B-12	NR	NR
Vitamin A	NR	NR
Vitamin E	NR	NR
Vitamin K	NR	NR
Carotene	7,807 µg	0
Cryptoxanthin	NR	NR
Lutein + Zeaxanthin	NR	NR

NR= not reported

Source: Puwastien et al., 2000

Review Article

Table 2: Constituents

Part	Constituents	Image
Fruits	Protien,fat,carbohydratesminerals:calcium,phosohorus,iron,vita min C, thiamine, riboflavin, niacin, carotene, Amino acid, citric acid and malic acid, voletile compound : linalool, benzylisothiocynate, cis and trans 2,6-dimethyl 3,6 epoxy-7 octen-2-ol,Alkaloid, α carpine, benzyl- β -D gucoside,2-phenyl ehyle- β -D gucoside,4-hydroxyphenyl-2-ehyle- β -D gucoside,	
Juice	N-butyric, n-hexanoic and n-octanoic acid, lipid, myristic, palmitic, stearic, linoleic, linolenic and cis-vasanic and oleic acid	
Seed	Fatty acid, crude protein, crude fibre, papaiya oil, carpaine, benzyl isothiocynate, benzyle glucosinolate, glucotropecolin, benzylethiourea, hentriaconate, β -sitosterol, caricin and enzyme myrosin	
Root	Carposide and enzyme myrosine	
Leaves	Alkaloids carpain, pseudocarpain and dhydrocarpaine 1 and 2 , choline, carposide, , vitamin C and E	
Bark	B-sitosterol, glucose, fructose, galactose and xylitol.	

Review Article

Latex Proteolytic enzyme, papain, chemopapain, glutamine cyclotransferase, chymopapain A, B, C, Peptidase A and B and lysozyme.



Source: Indian Material Medical by KM Nadkarni.
 Photochemistry PF Medicinal Plant, 2nd Edn by Jean Bruneton

Table 3: The cytotoxic effect of *C. papaya* extract has been tested in various cancer cell lines in vitro studies summarized intable

Cancer Cell Line	Treatment	Result	Reference
Breast cancer cell line (MDA-MB-231) Liver Cancer cell line (Hep G2) Chang liver cell line (normal cell)	Papaya fruit juice(0.28– 28 mg/mL), Lycopene extracted from papaya juice, Pure lycopene (3– 30 _g/mL)	Pure lycopene and papaya juice inhibited viability of liver cancer cell line Hep G2 (IC50 = 22.8 _g/mL and 20 mg/mL, respectively) but had no effect on breast cancer cells or normal cells. Lycopene extracted from papaya juice did not show any effect on either cell line.	Puwastien <i>et al.</i> , 2000
Acute promyelotic leukemia HL-60 cells	<i>n</i> -hexane extract of papaya seed or pulp (0.1– 100 _g/mL), Pure benzyl isothiocyanate (10 _M)	Extract of seed: Dose dependently inhibited the superoxide generation (IC50 = 10 _g/mL) and the viability of cells (IC50 = 20 _g/mL), comparable to that of pure benzyl isothiocyanate. Extract pulp had no effects at 100g/mL	Nakamura <i>et al.</i> , 2006
Breast cancer cell line (MCF-7)	Aqueous extract of papaya flesh (0.01–4% v/v)	Significant inhibitory effect on proliferation of MCF-7 cells ($p < 0.05$)	Garcia-Solis, 2009
Breast cancer cell line (MCF-7) treated with sodium nitroprusside, a nitric oxide donor	Ethanollic extract of papaya pericarp (50– 640 _g/mL)	Inhibited cell growth in MCF-7 cells (decrease in cell viability). Scavenged nitric oxide in dose-dependent manner (about 35% of nitric oxide was scavenged by extract at 640 _g/mL)	Jayakumar <i>et al.</i> , 2011
Breast cancer cell line (T47D)	Protein fraction containing RIPs isolated from leaves	The protein fraction possessed cytotoxicity: IC50 = 2.8 mg/mL). Induction of apoptosis by regulation of p53 and BCl-2 protein expression (increased by 59.4% and decreased by 63%, respectively).	Hirose <i>et al.</i> , 1998
Stomach cancer cell line (AGS) Panc0reatic cancer cell line	Aqueous extract of papaya	Papaya leaf extract showed a concentration-dependent anticancer effect on each of the cancer cell lines	Marotto <i>et al.</i> , 2006

Review Article

(Capan-1) Colon cancer cell line (DLD-1) Ovarian cancer cell line (Dov-13) Lymphoma cell line (Karpas) Breast cancer cell line (MCF-7) Neuroblastoma cell line (T98G) Uterine cancer cell line (Hela) T-cell leukemia cell line (CD26 negative or negative Jurkat)	leaves(1.25–27 mg/mL)	and suppressed DNA synthesis by suppressing the incorporation of 3H-thymidine.	
T-cell lines (H9, Jurkat, Molt-4, CCRF-CEM and HPB-ALL) Burkitt's lymphoma cell lines (Ramos and Raji) Chronic myelogenous leukemia cell line (K562) Cervical carcinoma cell line (Hela) Hepatocellular carcinoma cell lines (HepG2 and Huh-7)	Aqueous extract of papaya leaves (0.625–20 mg/mL)	Inhibited the proliferative responses of both haematopoietic cell lines and solid tumor cell lines. In peripheral blood mononuclear cells, papaya extract reduced the production of IL-2 and IL-4 whereas increased the production of Th1 type cytokines such as IL-12p40, IL-12p70, INF- γ , and TNF- α . The expression of 23 immunomodulatory genes was enhanced by the addition of papaya extract.	Otsuki <i>et al.</i> , 2010

Phytochemicals in c. Papaya with Reported Anticancer Activities

Carica papaya contains a broad spectrum of Phytochemicals including enzymes (in the latex), Carotenoids (in fruits and seeds), alkaloids (in leaves), phenolics (in fruits, leaves, and shoots), glucosinolate (in seeds and fruits) (Krishna *et al.*, 2008; Parle *et al.*, 2011). Some important Phytochemicals found in *C. papaya* are Lycopene, Betacarotinoide, Benzylisothiocyanate, Betacryptoxanthin, Benzylglucosinolate, chlorogenic acid, caffeic acid, protocatechuic acid, Quercetin, Among more than 5000 compounds from plants that have been identified to be associated with anticancer properties (Huang *et al.*, 2009), three groups of bioactive compounds-phenolics, Carotenoids, and glucosinolate-have attracted considerable interest in anticancer studies. Pure compounds of these three groups have been extensively researched in vivo and in vitro studies on many types of cell lines for their potential effects in cancer treatment and prevention. These bio actives act via multiple mechanisms such as cancer cell signalling, proliferation, apoptosis, migration, invasion, as well as angiogenesis and carcinogen elimination (Huang *et al.*, 2009; Zhang *et al.*, 2004; Thornalley *et al.*, 2002; Nakamura *et al.*, 2006; Wu *et al.*, 2009; Navarro *et al.*, 2011; Wahle *et al.*, 2011; Soobrattee *et al.*, 2006; Tanaka *et al.*, 2012; Breemen *et al.*, 2008) to exhibit in vitro and in vivo anticancer activities.

Papaya is Rich Source of Enzyme Papain which Effective against Cancer

Papain (EC 3.4.22.2) is an endolytic plant cysteine protease enzyme which is isolated from papaya (*Caricapapaya* L.) latex.(Abu-Alruz *et al.*, 2009)It preferentially cleaves peptide bonds involving basic amino acids, particularly arginine, lysine and residues following phenylalanine. (Menard *et al.*, 1990) The unique structure of papain gives its functionality that helps to understand how this Proteolytic enzyme works and it's useful for a variety of purposes. (*Caricapapaya* L.) Latex, (Mitchel, 1970).

Review Article



Figure 1: Papain Structure (Huet J *et al.*, 2006)

Many cancer cells having a protective coating of fibrin. That is why they go undetected for many months and years.

Papain breaks down that fibrin coat of cancer cell wall. So ultimately it helps against the cancer.

Mechanism of Functions of Papain

The mechanism in which the function of papain is made possible is through the cysteine-25 portion of the triad in the active site that attacks the carbonyl carbon in the backbone of the peptide chain freeing the amino terminal portion. As this occurs throughout the peptide chains of the protein, the protein breaks apart. The mechanism by which it breaks peptide bonds involves deprotonation of Cys-25 by His-159. Asparagine-175 helps to orient the imidazole ring of His-159 to allow this deprotonation to take place. Although far apart within the chain, these three amino acids are in close proximity due to the folding structure. It is through these three amino acids working together in the active site that provides this enzyme with its unique functions. Cys-25 then performs a nucleophilic attack on the carbonyl carbon of a peptide backbone. (Menard *et al.*, 1990; Tsuge *et al.*, 1999) In the active site of papain, Cys -25 and His -159 are thought to be catalytically active as a thiolate imidazolium ion pair. Papain can be efficiently inhibited by peptidyl or non-peptidyl N-nitrosoanilines. (Guo *et al.*, 1996; 1998) The inactivation is due to the formation of a stable S-NO bond in the active site (S-nitroso- Cys25) of papain. (Xian *et al.*, 2000)

Papaya is a Store-House of Cancer Fighting Lycopene

Lycopene is a member of the carotenoid family, which is synthesized by many plants and microorganisms. It is a highly unsaturated open straight chain hydrocarbon compound consisting of 11 conjugated and 2 unconjugated double bonds (Stahl *et al.*, 1992; Knachik *et al.*, 2002; Rao *et al.*, 2006). The red color of many fruits and vegetables is due to the presence of lycopene. Because of the presence of double bonds in the structure of lycopene, it can exist in both the *cis* and *Trans* isomeric forms. Lycopene is present in foods primarily in the all-*trans* isomeric form, (Cliton, 1998) However, it can undergo mono- or poly-isomerization by light, thermal energy, and chemical reactions to the *cis* isomeric form. It is highly stable at high temperatures and can be stored. (Agrawal *et al.*, 2001).




Figure 2: Structure of lycopene (Gali *et al.*, 2002)

Review Article

Mechanism of Lycopene

Lycopene enhances cell-to-cell communication by increasing 'Gap junctions' between cells (Gali-Muhtasib *et al.*, 2002). Two major mechanisms have been proposed to explain the ant carcinogenic activities of lycopene: non oxidative and oxidative mechanisms. Lycopene is hypothesized to suppress carcinogen-induced phosphorylation of regulatory proteins such as p53 and Rb antioncogenes and to stop cell division at the G0-G1 cell cycle phase (Obermuller-Jevis *et al.*, 2003). Astorg and colleagues proposed that lycopene-induced modulation of the liver metabolizing enzyme cytochrome P4502E1 was the underlying mechanism of its protection against carcinogen- induced preneoplastic lesions in rat liver. Preliminary *in vitro* evidence also indicates that lycopene reduces cellular proliferation induced by insulin-like growth factors-which are potent mitogens-in various cancer cell lines (Levy, 1995). Regulation of intrathymic T-cell differentiation (immune modulation) was suggested to be the mechanism for suppression of mammary tumor growth by lycopene treatments in SHN retired mice (Nagasava *et al.*, 1995 and Kobayashi *et al.*, 1996). This is an important mechanism by which cells communicate with each other, and helps to ensure proper cell and organ functions. It also prevents uncontrolled growth of cancer cells and modulates cell-cycle progression. The cell cycle is a highly ordered set of events that culminates in cell division. The use of lycopene appears to be a potential anticancer strategy of regulating the cell cycle by inhibiting abnormal cellular growth (Agrawal *et al.*, 1998). Lycopene has been hypothesized to prevent carcinogenesis by protecting critical cellular bimolecular, including lipids, lipoproteins, proteins, and DNA (Matsushima *et al.*, 1995; Pool-Zobel *et al.*, 1997; Rao *et al.*, 1998).

Isothiocyanate Found in Papaya Restore the Cell Cycle to Eliminate Cancer

Organo-sulfur compounds called isothiocyanate are found in papaya. In animal experiments, isothiocyanate protected against cancers of the breast, lung, colon pancreas, and prostate, as well as leukemia, and they have the potential to prevent cancer in humans. Isothiocyanate have shown them capable of inhibiting both the formation and development of cancer cells through multiple pathways and mechanisms (Barbara, October-2008).

Researchers in Japan clarified the mechanisms of action in a type of isothiocyanate found in papaya known as BITC. That underlies the relationship between cell cycle regulation and appropriate cell death. When cancerous cells die on schedule, they are no longer a problem. The researchers established that BITC exerted cancer cell killing effects that were greater in the proliferating cells than in the quiescent cells. Cancer cells that are proliferating are much more dangerous than cancer cells that are in a state of dormancy (Barbara, 2009).

CONCLUSION

Papaya is originally native to southern Mexico and now cultivated in many tropical counties. Sweet and succulent with satiny consistency. Papaya was referring to as the fruit of the angels. Slice open a papaya and see hundreds of shiny black seeds that all need to get there start in life from the nutrition found in the fruit. This implies that fruit must be power packed. The papaya plant has been touted by traditional hearlers for counties as source of powerful medicine. Papaya not only the delicious fruit but it is loaded with vitamins, Phytochemicals, and several mineral compound and hawing some important enzyme like papain, lycopene, Isothyocynate and some proteolysis enzyme which help to treat health problem. Papaya promotes immune system papaya is potent cancer fighter that is highly effective against hormone related to cancer as well as other cancer. Papaya can stop the growth of cancer cell halt metastasis and normalized cell cycle.

REFERENCES

Abu-Alruz K, Mazahreh AS, Quasem JM, Hejazin RK and El-Qudah JM (2009). Effect of Proteases on Melt ability and Stretch ability of Nabulsi Cheese. *American Journal of Agriculture and Biology Science* **4** 173-178.

Review Article

- Agarwal A, Shen H, Agarwal S and Rao A (2001).** Lycopene content of tomato products, its stability, bioavailability, and in vivo antioxidant properties. *Journal of Medicinal Food* 15-49.
- Agarwal S and Rao AV (1998).** Tomato lycopene and low density lipoprotein oxidation: A human dietary intervention study. *Lipids* 33 981-984.
- Barbra L Minton (2008).** Papaya is tasty way to fight cancer and poor digestion. *International Journal of Oncology*.
- Barbra L Minton (2009).** Papaya is tasty way to fight cancer and poor digestion Forum of nutrition.
- Bhattacharjee SK (2001).** Carica papaya. In: *Hand Book of Medicinal Plant*, edition: 3rd Revised, editors: Shashi Jain, (Pointer Publisher, Jaipur) 1-71.
- Clinton SK (1998).** Lycopene: Chemistry, biology and implications for human health and diseases. *Nutrition Reviews* 1 35-51.
- Gali-Muhtasib H and Bakkar N (2002).** Modulating cell cycle: Current applications and prospects for future drug development. *Current Cancer Drug Targets* 2 309-336.
- Garcia-Solis P, Yahia, EM, Morales-Tlalpan V and Diaz- Munoz M (2009).** Screening of ant proliferative effect of aqueous extracts of plant foods consumed in Mexico on the breast cancer cell line MCF-7. *International Journal of Food Sciences and Nutrition* 60 32-46.
- Ghosh S (2005).** Physicochemical and conformational studies of papain/sodium dodecyl sulfate system in aqueous medium. *Journal of Colloids and Surfaces A: Physicochemical and Engineering Aspects* 264 6-16.
- Hirose M, Yamaguchi T, Kimoto N and Ogawa K (1998).** Strong promoting activity of phenyl ethyl isothiocyanate and benzyl isothiocyanate on urinary bladder carcinogenesis in F344 male rats. *International Journal of Cancer* 77 773-777.
- Huang WY, Cai YZ and Zhang Y (2009).** Natural phenolics compounds from medicinal herbs and dietary plants: potential use for cancer prevention. *Nutrition and Cancer* 62 1-20.
- Jayakumar R and Kanthimathi MS (2011).** Inhibitory effects of fruit extracts on nitric oxide-induced proliferation in MCF- 7 cells. *Food Chemistry* 126 956-960.
- Jean Bruneton (1999).** Carica Papaya. In: *Pharmacognosy, Photochemistry of Medicinal Plant*, edition: 2nd, Technique and Documentation, France 221-223.
- Knachik F, Caryallo L, Bernstein PS, Muir GJ, Zhao DY and Katz NB (2002).** Chemistry, distribution and metabolism of tomato Carotenoids and their impact on human health. *Experimental Medicine and Biology* 227 845-851.
- Kobayashi T, Iijima K, Mitamura T, Torizuka K, Cyong JC and Nagasawa H (1996).** Effects of lycopene, a carotenoid, on intrathymic T cell differentiation and peripheral CD4/CD8 ratio in a high mammary tumor strain of SHN retired mice. *Anticancer Drugs* 7 195-198.
- Krishna KL, Paridhavi M and Patel JA (2008).** Review on nutritional, medicinal and pharmacological properties of papaya (*Carica papaya* Linn.). *Natural Product Radiance* 7 364-373.
- Levy J, Bosin E, Feldmen B, Giat Y, Miinster A and Danilenko M (1995).** Lycopene is a more potent inhibitor of human cancer cell proliferation than either α -carotene or β -carotene. *Nutrition and Cancer* 24 257-266.
- Marotta F, Weksler M, Naito Y, Yoshida C, Yoshioka M and Marandola P (2006).** Nutraceutical supplementation, effect of a fermented papaya preparation on redox status and DNA damage in Healthy elderly individuals and relationships with GSTM1 genotype, a randomized, placebocontrolled, cross-over study. *Annals of the New York Academy of Sciences* 1067(1) 400-407.
- Matsushima NR, Shidoji Y, Nishiwaki S, Yamada T, Moriwaki H and Muto Y (1995).** Suppression by Carotenoids of microcrystal-induced morphological changes in mouse hepatocytes. *Lipids* 30 1029-1034.
- Menard R, Khouri HE, Plouffe C, Dupras R and Ripoll D (1990).** A protein engineering study of the role of aspartate 158 in the catalytic mechanism of papain. *Biochemistry*, 29 6706-6713.
- Mike Barrett (2012).** Carica Papaya a Powerful Cancer Fighter.

Review Article

- Mitchel RE, Claiken MI and Smith ELJ (1970).** The complete amino acid sequence of papain. *The Journal of Biological Chemistry*.
- Nagasawa H, Mitamura T Sakamoto S and Yamamoto K (1995).** Effects of lycopene on spontaneous mammary tumour development in SHN virgin mice. *Anti Cancer Research* **15** 1173-1178.
- Nakamura Y and Miyoshi N (2006).** Cell death induction by isothiocyanate and their underlying molecular mechanisms. *Bio Factors* **26** 123-134.
- Nakamura Y, Yoshimoto M, Murata Y and Shimoishi Y (2007).** Papaya seed represents a rich source of biologically active isothiocyanate. *Journal of Agricultural and Food Chemistry* **55** 4407-4413.
- Navarro SL, Li F and Lampe JW (2011).** Mechanisms of action of isothiocyanate in cancer chemoprevention: an update. *Journal of Food Function* **2** 579-587.
- Obermuller-Jevis UC, Olano-Martin E, Corbacho AM, Eiserich JP, Van Der Vliet A and Valacchi G (2003).** Lycopene inhibits the growth of normal human prostate epithelial cells in vitro. *Journal of Nutrition* **133** 3356-3360.
- Otsuki N, Dang NH, Kumagai E and Kondo A (2010).** Aqueous extract of Carica papaya leaves exhibits anti-tumor activity and immunomodulatory effects. *Journal of Ethno Pharmacology* **127** 760-767.
- Parle M and Gurditta (2011).** Basketful benefits of papaya. *International Research Journal of Pharmacy* **2** 6-12.
- Pool-Zobel BL, Bub A, Muller H, Wollowski I and Rechkemmer G (1997).** Consumption of vegetables reduces genetic damage in humans: First result of a human intervention trial with carotenoid-rich foods. *Carcinogenesis* **18** 1847-1850.
- Puwastien P, Burlingame B, Raroengwicht M and Sungpuag P (2000).** *ASEAN Food Composition Tables of Nutrition*, (Mahidol University, Thailand).
- Rahmat A, Rosli R, Endrini S and Zain SAH (2002).** Ant proliferative activity of pure lycopene compared to both extracted lycopene and juices from watermelon (*Citrullusvulgaris*) and papaya (*Carica papaya*) on human breast and liver cancer cell lines. *Journal of Medical Science* **2** 55-58.
- Rao AV and Agarwal S (1998).** Bioavailability and in vivo antioxidant properties of lycopene from tomato products and their possible role in the prevention of cancer. *Nutrition Cancer* **31** 199-203.
- Rao AV, Mira MR and Rao LG (2006).** Lycopene. *Advances in Food and Nutrition Research* **51** 99-164.
- Soobrattee MA, Bahorun T and Aruoma OI (2006).** Chemo preventive actions of polyphenolic compounds in cancer. *Biomedical Factors* **27** 19-35.
- Stahl W, Schwarz W, Sundquist AR and Sies H (1992).** Cis-Trans, Isomer of Lycopene and Beta-carotene in human serum and tissues. *Archives of Biochemistry and Biophysics* **294** 173-177.
- Tanaka T, Shnimizu M and Moriwaki H (2012).** Cancer chemoprevention by Carotenoids. *Molecules* **17** 3202-3242.
- Thornalley PJ (2002).** Isothiocyanate: mechanism of cancer Chemo preventive action. *Anticancer Drugs* **13** 331-338.
- Tsuge H, Nishimura T, Tada Y, Asao T and Turk D (1999).** Inhibition mechanism of cathepsin Lspecific inhibitors based on the crystal structure of papain-CLIK148 complex. *Journal of Biochemical and Biophysical Research Communications* **266** 411-416.
- Van Breemen RB and Pajkovic N (2008).** Multitargeted therapy of cancer by lycopene. *Cancer Letters* **269** 339-351.
- Wahle KWJ, Brown I, Rotondo D and Heys SD (2011).** Plant phenolics in the prevention and treatment of cancer. *Bio-Farms for NutraSweet* **698** 36-51.
- Wu X, Zhou QH and Xu K (2009).** Are isothiocyanate potential anti-cancer drugs? *Acta Pharmacological Sinica* **30** 501-512.
- Xian M, Chen X, Liu Z, Wang K and Wang PG (2000).** Inhibition of papain by *s*-nitrosothiols. *Journal of Biological Chemistry* **275** 20467-20473.

Review Article

Zhang Y (2004). Cancer-preventive isothiocyanate: measurement of human exposure and mechanism of action. *Mutation Research: Fundamental and Molecular Mechanisms* **555** 173-190.